

The detection and determination of diene hydrocarbons with a conjugated system of double bonds. I. V. I. Kaslov, *J. Applied Chem.* (U. S. S. R.) 14, 141-47 (in German), 1471 (1941); translated in *Foreign Petroleum Tech.* 9, 443-50 (1941). The bromometric Kaufmann method (C. A. 30, 7885⁵) cannot be used universally for the detection of diene hydrocarbons with conjugated double bonds. The excessively low values of the polymerization of dienes are due to side reactions of the polymers with monomers by Diene hydrocarbons are distinguished from monoalkenes by the generation of considerable amounts of HBr when brominated in a soln. of CCl_4 . The Mellin method is suitable for detection of dienes alone or in mixts. of various hydrocarbons. II. V. I. Kaslov and A. V. Shpash, *J. Applied Chem.* (U. S. S. R.) 14, 148-50 (in German), 1501 (1941); translated in *Foreign Petroleum Tech.* 9, 350-51 (1941).—Bromination in CCl_4 is specific for diene hydrocarbons with conjugated double bonds and is accompanied by the generation of considerable amounts of halogen acids. Hydrocarbon chains with separated double bonds are brominated similarly to olefins, with an insignificant amount of halogen acid. The sensitivity limits of the bromination reaction for mixts. of olefins with 1,3-dienes is 1% of the latter, while for mixts. of satd. hydrocarbons with dienes this limit is at 0.5% of dienes. The experiments were carried out with linseed oil and mixts. of cyclohexene and decalin, and satd. kerosene and decalene, by the Mellin method of detg. I nov. A. A. Bochtling

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962920002-0"

COMMON ELEMENTS		RARE EARTH AND METALS		FACILITIES AND PROPERTIES INDEX		OTHER SYMBOLS					
COMMON ELEMENTS		RARE EARTH AND METALS		FACILITIES AND PROPERTIES INDEX		OTHER SYMBOLS					
<p><i>ca</i></p>		<p><i>10</i></p>		<p>Characteristics of α,β-unsaturated ketones. II. V. F. Kozlov and I. I. Ralibier. J. Gen. Chem. (U.S.S.R.) 13, 209-217(1943) (English summary); cf. C. A. 13, 3056.</p> <p>It was shown that the displacement of ketones by aldehyde from α,β-unsat. ketones is connected with the preliminary hydrolytic cleavage of the latter into its components. BzH₂, treated with mesityl oxide in the presence of dil. NaOH, yielded small amounts of benzylidenacetone and benzylidenmesityl oxide, the latter being isolated as the tetrahydrate, m. 118° (from EtOH). The reactions were conducted by prolonged standing at about 10°, with a longer period noticeably increasing the yield of benzylidenacetone. Me₂CO was shown to be inert to condensation on treatment with piperidine acetate on prolonged heating on a steam bath; the same result was obtained when MeEtCO was used. However, 25 g. BzH₂, 30 g. Me₂CO, 5 g. piperidine and 5 g. AcOH yielded, after 20 hrs. heating, 71% benzylidenacetone, m. 41.8°, while similar reaction with mesityl oxide gave crude benzylidenmesityl oxide bp. 173-5°, identified as the tetrahydrate. Thus, piperidine acetate may be recommended for aldehydic-ketone condensations.</p> <p style="text-align: right;">G. M. Kovalevich</p>		<p>ABN-SLA METALLURGICAL LITERATURE CLASSIFICATION</p>		<p>FROM SYNONYM</p>		<p>FROM COMMON</p>	
<p>SYMBOL NO.</p>		<p>SERIES WIP ONE UNIT</p>		<p>CALCULATION</p>		<p>WEIGHT OF CHG. IN</p>					
<p>A B C D E F G H I J K L M N O P Q R S T U V W X Y Z</p>		<p>A B C D E F G H I J K L M N O P Q R S T U V W X Y Z</p>		<p>A B C D E F G H I J K L M N O P Q R S T U V W X Y Z</p>		<p>A B C D E F G H I J K L M N O P Q R S T U V W X Y Z</p>					

PROCEDURES AND PROPERTIES INDEX	
<p>10</p> <p>Preparation of enanthylidenacetone. V. I. Kuznetsov, I. P. Vladimirov, M. S. Kasakhina, Z. S. Elida, Z. I. Pronina and I. I. Ralshar. <i>J. Gen. Chem.</i> (U.S.S.R.) 13, 814-17 (1943) (English summary).—It was shown that the product obtained by Rupe and Hinterlach (C. A. J. 800) from condensation of MeCO with enanthole in the presence of 1% NaOH is not enanthylidenacetone but decen-4-ol-5-one. Authentic enanthylidenacetone was prepd. as follows: 5 g. piperidine and 3 g. ethyl AcOH, in 50 g. dry MeCO, were treated dropwise with 20 g. enanthole while refluxing the mixt. on a steam bath; after the addn., which requires 4 hrs., the mixt. was heated for 8 hrs., cooled, the aq. layer sepd. while the org. layer was freed of MeCO, dild. with EtO, dried over Na₂SO₄ and distd. to yield enanthylidenacetone, bp 100-102°, n_D²⁰ 1.4480, d₄²⁰ 0.8473; yield, 15-17 g. The product yields the normal positive iodoform reaction. The product obtained by R. and H. yields enanthylidenacetone after dehydration over KHSO₄-Na₂SO₄ by repeated distn. at 22 mm.</p> <p>G. M. Kouslapoff</p>	
<p>ASS-SLA METALLURGICAL LITERATURE CLASSIFICATION</p>	
<p>FROM STUDY</p>	
<p>FROM WORK</p>	

10

Ca

Synthesis of new hydrocarbons with a conjugated system of double bonds. IV. V. A. Egorov, I. P. Chudakov, M. S. Kasilkina and I. I. Nalcher. *J. Gen. Chem.* (U. S. S. R.) 13, 818-22 (1943) (English summary); cf. C. A. 39, 476^a and following abstr.—MeMgl (from 14.5 g. MeI) was added dropwise over 3 hrs. to 13.4 g. cyanideacetone in Et₂O at -15°; after standing overnight and decomposition by ice and NH₄Cl, there was obtained 16% 3-methyl-1,3-decadiene, bp 88°, n_D^{20} 1.4422, d_4^{20} 0.7718. The product, heated for 20 hrs. in a steam bath in a sealed tube with maleic anhydride in toluene, yielded an *adduct*, apparently of the trimer, which could not be reduced to the crystalline state. Similar reaction using EtMgBr gave a mixt. of 3-methyl-2,6-hexadiene and 3-ethyl-1,3-octadiene, bp 96-104°, n_D^{20} 1.4510, d_4^{20} 0.7903; heating with maleic anhydride gives a mixt. of adducts of the monomer and the dimer of the hydrocarbons; in the form of a heavy oil. A similar prepn. using *iso*-AmMgBr gave a mixt. of 3-isopropyl-1,3-decadiene and 3,6-dimethyl-1,3-octadiene, bp 120-34°, n_D^{20} 1.4164, d_4^{20} 0.7866. G. M. Kostikov

ASS-514 METALLURGICAL LITERATURE CLASSIFICATION

SEARCHED INDEXED

RECEIVED

LIBRARY

DATE

BY

NO.

10

1ST AND 2ND EDITIONS

SYNOPSIS AND PROPERTIES INDEX

7

CA

Detection of diene hydrocarbons with a conjugated double-bond system. III. V. I. Kozlov, J. Appl. Chem. (U. S. S. R.) 16, 283-84 (1943) (English summary); J. C. A. 16, 2223. On the basis of more accurate work it is shown that the reaction of bromination in CCl_4 is specific only for those dienes which have 1, or better 2, side chains connected to carbons of the conjugated system; in the latter case considerable HBr is evolved and 80-90% of Br_2 is used for these side reactions. Di- and tri-isobutylenes are characterized by this behavior among the olefin hydrocarbons. The bromination reaction is specific for dienes of normal and branched structures and these are characterized by some 80% conversion of I into III by the side reactions; di- and tri-isobutylenes are characterized likewise. The I an. procedure of H. G. Waller is most precise for detg. analyt. Q. H. Kozlov

ABB-55.5 METALLURGICAL LITERATURE CLASSIFICATION

CODE SYMBOLS

1ST AND 2ND EDITIONS

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

CR

10

Effecting the Grignard reaction with α,β -unsaturated ketones. V. V. I. Kuznetsov. *J. Gen. Chem. (U.S.S.R.)* 14, 84-7 (1944) (English summary); cf. preceding abstr. ... Simultaneous addn. of EtMgBr (in Et_2O) and methyl oxide (in Et_2O) to Et_2O kept at -10° , followed by the usual working-up and dehydration of the crude tertiary alk. with $\text{CaH}_2(\text{CO})_2$, gave an av. yield of 50.9% 3,4-dimethyl-7,8-octadiene, b. $114-15^\circ$, thus showing that, in reactions of Grignard reagents with α,β -unsatd. ketones, the diln. and employment of low temp. favors the normal reaction course. The same reaction, conducted at 15° , gave 17.5% of crude octadienes, consisting of 3-methyl-4-methyl-1,3-pentadiene, b. $111-12^\circ$, d_4^{20} 0.7533, n_D^{20} 1.4424, and an undetd. amt. of the higher-boiling isomer. The product of the -10° reaction was sepd., in the ratio of 1:1, into the above pentadiene and the hexadiene, b. $114-15^\circ$, d_4^{20} 0.7542, n_D^{20} 1.4438. The structures of the compounds were detd. by oxidation. G. M. Kosolapoff

117 AND 118 INDEX		PROCESSING AND PROPERTY INDEX		119 AND 120 INDEX	
<p>ca</p> <p>10</p> <p>Attempted synthesis of a ketone of the cyclopenteno- arben. I. V. I. Kozlov. <i>J. Gen. Chem. (U.S.S.R.)</i> 14, 200-202 (1964) (English summary).--Attempted cycliza- tion between 3-methyl-5-ethyl-2,6-heptadiene and PhNC failed because of the ease of dimerization of the latter. The expts. were conducted in sealed tubes at room temp for prolonged periods of time and with heating on a steam bath as well as on an oil bath at 133-135°. Phenyl iso- nitrile dimer is a cryst. solid, m. 135° (from petr. ether); it is rather readily hydrolyzed, thus indicating an imidic linkage between C and N, and it probably has the structure</p> $\text{PhN}=\text{C} \quad \text{PhN}=\text{C}$ $\text{C}=\text{NPh} \quad \text{or} \quad \text{C}=\text{NPh}$ <p>G. M. Kozlov</p>					
<p>ASACLA METALLURGICAL LITERATURE CLASSIFICATION</p> <p>117 AND 118 INDEX</p> <p>119 AND 120 INDEX</p>					

Handwritten: 10

PROCEDURES AND RECOVERY INDEX

Halogenation of unsaturated hydrocarbons. IV. V.
I. Eskin (Sverdlovsk State Univ.). J. Gen Chem,
(U.S.S.R.) 15, 1001-6(1945); cf. C.A. 39, 470*.—The
Macleaney method of iodine no. detn. is recommended for
detection of structures at the quaternary C atom having a
double bond, such as in olefins or in dienes. The amt. of
such hydrocarbons in liquid-phase cracking products is
shown to increase with rising b.p. of the fractions. A
hypothesis for halogenation and hypohalogenation of
olefins is proposed according to which the halogenating
mol. is first fixed to the double bond, then the atomic
bonds are cleaved, and the halogens are transformed into
stains nascendi; this explains the possibility and the high
reaction rate of such halogens with other materials present
in the fraction mixt. Isoamylene (1.00 g.) and 8.9922 g.
kerosene fraction with iodine no. 0.0 were mixed and 4.7085
g. of the edn. was treated for 0.5 hr. with 2.5 l. freshly
precip. Cl water, and the org. layer was sepl., washed with
dil soda soln., dried, and analyzed for Cl and unreact.;
19% of the isoamykene did not enter into reaction, thus
showing direct chlorination of isoamykene; extension of
the reaction to 1 hr. gave 17% unreacted isoamykene.
The supposition that hypochlorination is accompanied by
direct chlorination is also supported by the higher Cl con-
tent than calcd. for an HOCl addn. product. Bromina-

tion of 2,4-dimethyl-2,4-hexadiene To CC1₄ at 0° for
0.5–2.0 hrs. showed that 8.8–29.2% of total utilized Br is
used for transformation into HBr. G. M. Kosoborod

Handwritten: 10

UNITED STATES GOVERNMENT PRINTING OFFICE: 1947 O - 210,000

1ST AND 2ND CODES		PROCESSIES AND PROPERTIES INDEX	
<p>Chemical composition of the fatty oil from <i>Lappula echinata</i> Glib. V. I. Easlov, V. L. Glizina, and M. A. Panyukova. <i>J. Applied Chem. (U.S.S.R.)</i> 18, 176-0 (1945).—The oil from seeds of <i>L. echinata</i> can be used in formulation of high-grade drying oils. Among the constituents there was found a new form of linolenic acid,</p>		<p>whose hexabromide, m. 232-4°, obtained by catn. of the crude hexabromides with ether. G. M. Kosolapoff</p>	
<p>ASD-51A METALLURGICAL LITERATURE CLASSIFICATION</p>			
<p>FROM LITERATURE</p>		<p>RELATIONS</p>	
<p>SEARCHED MAY ONE ONE</p>		<p>SEARCHED ONE ONE ONE</p>	

Synthesis of new diene hydrocarbons with a conjugated system of double bonds. VI. V. I. Ruzhkov and V. V. Mikhlinova (Ural State Univ., Sverdlovsk). *J. Gen. Chem.* (U.S.S.R.) 16, 1883 (1946); cf. *C.A.* 39, 1919. With the improved Grignard technique (*C.A.* 39, 1919) involving primarily low temp., high order of dist., and slow addition of the Grignard reagent, a no. of diene hydrocarbons were prepared in satisfactory yields from α,β -unsatd. ketones. Generally, ketones of higher mol. wt. give lower yields, and Me ketones react less readily than Et ketones, as far as the normal reaction is concerned. Aromatic Grignard reagents favor abnormal course of the reaction to a greater extent than Me or Et Grignard reagents, in contradiction to Kohler (*C.A.* 3, 124). *EtMgBr* (from 3.8 g. *Me* and 21 g. *EtBr*) in 40 cc. *Et₂O* was added over 6 hrs. at -14° to 20 g. 3-methyl-3-hepten-5-one in 100 cc. *Et₂O*. After working up as described in *C.A.* 33, 3958, 82.9% 3-methyl-5-ethyl-3,5-heptadiene, b.p. 154° , d_4^{20} 0.7710, n_D^{20} 1.4456 was obtained. To 41 g. 4-octen-3-one, b.p. $171-3^{\circ}$, d_4^{20} 0.8541, n_D^{20} 1.4653, in 130 cc. *Et₂O*, cooled to -15° , *EtMgBr* (from 11.5 g. *Me* and 60 g. *EtBr*) in 80 cc. *Et₂O* was added over 6 hrs; after standing overnight and decan., with ice and *NH₄Cl*, the dried *Et₂O* ext., treated with hydroquinone, distd. in *CO₂* over 2 g. *KHSO₄* and 5 g. anhyd. *Na₂SO₄*, and finally purified by repeated distn. over *Na*, gave 26.6% 3-ethyl-3,5-octadiene, b.p. 167° , d_4^{20} 0.7727, n_D^{20} 1.4620; the hydrocarbon (3 g.) and 3.2 g. maleic anhydride, heated in 10 cc. *Me₂Ph* in a sealed tube for 100 hrs., gave an acid forming an Ag salt with 34.74% Ag.

which appears to correspond most closely to an adduct of the hydrocarbon dimer; inspection of the condensation at room temp. with 15 min. heating to 65° gave an oily product contg. 37.71% Ag; finally, the diene (3 g.) heated with 2.4 g. quinone in 10 cc. *Me₂Ph* to 100° for 2 hrs., gave 0.9 g. *Quinone*, m. 132° , which readily reacts with *Et* in *CCl₄*. To 10.5 g. *PrCH:CHCOMe*, b.p. $164-7^{\circ}$, d_4^{20} 0.8114, n_D^{20} 1.4417, in 100 cc. *Et₂O* cooled to -15° to -20° , *MeMgI* (from 2.3 g. *Me* and 17 g. *MeI*) in 30 cc. *Et₂O* was added over 6 hrs.; treatment as above gave 16% 3-methyl-1,3-heptadiene, b.p. $127-8^{\circ}$, d_4^{20} 0.7453, n_D^{20} 1.4632, while a doubled amt. of *MeMgI* gave 20.5% the Diels-Alder adduct with maleic anhydride to the adduct Ag salt, contg. 35.95% Ag (corresponding to the adduct of the dimer or trimer). To 17.0 g. *PrCH:CHCOMe* in 100 cc. *Et₂O*, cooled to -10° to -15° , *PhMgBr* (from 6 g. *Me* and 34 g. *PhBr*) in 80 cc. *Et₂O* was added over 6 hrs. to yield, upon dehydration as above, but at reduced pressure, and vacuum distn. over *Na*, and finally atm. distn. over *Na*, 2.2 g. 3-phenyl-1,3-heptadiene, b.p. $234-7^{\circ}$, d_4^{20} 0.9553, n_D^{20} 1.5430; the maleic anhydride adduct (prepd. at 120°) gave an Ag salt, contg. 40.97% Ag, which was fairly close to *C₁₂H₁₆O₄*. G. M. K.

Aluminum chloride-catalyzed addition of *tert*-butyl chloride to propylene. Verle A. Miller (General Motors Corp., Detroit, Mich.). *J. Am. Chem. Soc.* 69, 1704-8 (1947). When 1.25 moles of *C₄H₈* is bubbled into 1 mole *tert*-BuCl (1) and 2 g. *AlCl₃* at -45° , at such a rate that the absorption of the gas is complete, 4-10% of the *C₄H₈* and 23-3% 1 are recovered and 70% of a butyl chloride

fraction (II) is formed. A curve shows the effect of reaction temp. and quantity of catalyst. About 45% of II is composed of 2-chloro-4,6-dimethylpentane; the remainder is principally 3-chloro-2,3-dimethylpentane with a small quantity of the 2-Cl isomer. These were identified by reaction with the Grignard reagent and comparison of the phys. properties and derivs. with known compounds. The principal fraction of the higher-boiling mixt. obtained is composed chiefly of isopropyl chlorides which were not further identified. At higher reaction temps. and when more catalyst was used, the reaction was more complex; the amt. of II decreased and the primary addition product was converted almost entirely into the rearranged product; the quantity of higher-boiling product obtained increased; the quantity of lower-boiling side-reaction products increased from a negligible to a significant quantity. 2,3-Dimethyl-3-pentanol yields a *phenylacetone*, m. 48.5-9°; that from 4,4-dimethyl-3-pentanol m. 52-3°.

C. J. West

27

B

Halogenation of the Unsaturated Hydrocarbon. V.
(In Russian.) V. I. Esafov, Zhurnal Obshchey Khimii
(Journal of General Chemistry), v. 17(70),
Aug. 1947, p. 1463-1467.

Presents results of an experimental and theoretical investigation of the bromination of mono- and diolefins of the aliphatic series and of the stability of the products.

METALLURGICAL LITERATURE CLASSIFICATION

ESSAFOV, V. I.

"On the Attempt to Synthesize the Ketones of the Cyclopentene Series. II"
(p. 1518)

SO: Journal of General Chemistry, (Zhurnal Obshchei Khimii), 1947, Vol. 17, No. 8

YESAFOR, V. L.

PA 64/49T19

USSR/Chemistry - Diene Synthesis Jun 49
Chemistry - Grignard Reaction

"Formulating and Isolating New Diene Hydrocarbons With Conjugate Double Bonds," V. L. Yesafor, Lab of Org Chem, Ural State U, Sverdlovsk, 14 pp

"Zhur Obshch Khim" Vol XIX, No 6 - p.1065

Gives syntheses of 2, 6-dimethylheptadiene-1, 3; 3-methyloctadiene-2, 4; 2-tertiary-butyl-4, 5, 5-trimethylheptadiene-1, 3; and 2, 3, 4-trimethylheptadiene-1, 3. Physicochemical constants of 2, 6-dimethyloctadiene-4, 6 are also more precise, and yield of 2, 4-

64/49T19

USSR/Chemistry - Diene Synthesis Jun 49
(contd)

dimethylpentadiene-1, 3 is improved. Studies Grignard reaction of three following systems:

- (1) 3-methylheptene-3-on-5 and C_6H_5MgI , (2) 3, 4-dimethylheptene-3-on-5 and C_6H_5MgBr , and (3) butylidene-methylpropylketone and C_6H_5MgBr . Describes tertiary alcohols, 2, 6-methyloctene-4-ol-6 and 3, 4, 5-trimethylheptene-4-ol-3. Shows that in the dehydration of α, β -unsaturated tertiary alcohols, having in the presence of carbinol the groups CH_3 and C_2H_5 , activity is chiefly caused by the H-atom of α -methyl. Submitted 9 Sep 47.

64/49T19

CA

The characteristics of α,β -unsaturated ketones. IV.
V. I. Esalov. *J. Gen. Chem. U.S.S.R.* 19, 1100-18
(1949) (Engl. translation).—See C.A. 44, 5793a. E. J. C.

CH

Characterization of α,β -unsaturated ketones. IV. V. I. Easlov, *Zhur. Obshchei Khim.* (J. Gen. Chem.) 19, 1115-24(1949); cf. *C.A.* 39, 918'.—Summarization of the previous data (*C.A.* 35, 3958'; 40, 6403'; 44, 10016) shows that the degree of conjugation of C:C and C:O links det. the reactivity of unsatd. ketones; increased mass of R bound to the CO group decreases the extent of conjugation, and the same takes place with the R attached to the terminal C of the C:C link. Two satd. radicals on the latter C atom decrease the conjugation even more, and the action of a Me-Et combination is greater than that of di-Me or *tert*-Bu. Introduction of a satd. R on the 2nd C atom, with 2 radicals on the 1st C atom, gives the max. lowering of the conjugation. The iodination of the unsatd. ketones may serve to show the structural types by increase of the iodine no. with increased no. of side chains from the C atoms of the conjugated

system; bromination in CCl_4 can also be used similarly, as the iodine no. of addn. decreases with an increased no. of radicals bound to the conjugated system. The upper conjugation limit is given by vinyl ketones and the lower limit lies with the compds. having 2 quaternary C atoms in the C:C link. The ketones used (the methods of prepn. will be given in later paper) were: butylidenemethyl ketone (I), b_{m} 163-6°, d_4^{20} 0.8520, n_D^{20} 1.4424; butylidenemethyl ethyl ketone (II), b_{m} 175-7°, d_4^{20} 0.8613, n_D^{20} 1.4470; butylidenemethyl propyl ketone (III), b_{m} 183-7°, d_4^{20} 0.8540, n_D^{20} 1.4450 (probably contains some 3-methyl-3-hepten-2-one); isomethylidenacetone (IV), b_{m} 100-10°, d_4^{20} 0.8473, n_D^{20} 1.4486; isomethylidenacetone (V), b_{m} 174-8°, d_4^{20} 0.8470, n_D^{20} 1.4409; mesityl oxide (VI), b_{m} 129-30°, d_4^{20} 0.8602, n_D^{20} 1.4440; 3-methyl-3-hepten-5-one (VII), b_{m} 164°, d_4^{20} 0.8591, n_D^{20} 1.4421; 3,4-dimethyl-3-hexen-5-one (VIII), b_{m} 157-8°, d_4^{20} 0.8585, n_D^{20} 1.4416; 3-methyl-4-methyl-3-hepten-2-one (IX), b_{m} 187-91°, d_4^{20} 0.8543, n_D^{20} 1.4450; 3,3,5,5-tetramethyl-4-hepten-3-one (X), b_{m} 202-3°, d_4^{20} 0.8525, n_D^{20} 1.4461; benzylidenemethyl ketone (XI), m 42.5°; and benzylidenemethyl ethyl ketone (XII), m 40°. The iodination was run by addn. of the ketone to 10 ml. EtOH, rinsing with 10 ml. EtOH, addn. of 25 ml. EtOH-iodine soln. and, after shaking, 20 ml. H_2O .

followed by shaking and letting stand 5 min. or longer, and titration of the excess iodine with thiosulfate, while HI is calcd. from the reaction with 10 ml. 2% KIO₃ and a 2nd titration with thiosulfate; brominations were done in CCl₄, the mixts. allowed to stand 30 min. after mixing, and the HBr generated calcd. similarly from the reaction with KI and KIO₃. In 5 min. I gave an iodine no. of 3.5-3.9 with 0.9-5.1% HI formation, while a 1370-min. reaction gave 27.5-32.2 and 79.1-85.5%, resp.; II in 5 min. gave 6.4-6.1 iodine no. and 26.3-32.4% HI, while 1330 min. gave 25.7-19.7 and 67.3-78.5%, resp.; III in 5 min. gave 34.5-5.2 and 10.5-18.0%, resp. (longer run not made); XI in 5 min. gave 5.1-3.0 iodine no. (HI not reported) while 1350 min. gave 15.1-18.6 and 15.3-17.0%, resp.; XII in 5 min. gave 8.8-5.9 and 36.5-7.0%, resp.; VI gave in 5 min. 3.2-6.5 iodine no. and no HI cited, while 1740 min. gave 105.4 and 46.9% HI, resp.; X in 5 min. gave 4.8-6.1 iodine no. and 35.1-44.0% HI; VII in 5 min. gave 36.6-4.5 iodine no. and 92.6-8.4% HI, while 1700 min. gave 156.7-3.5 and 59.2-64.0%, resp.; VIII in 5 min. gave 93.3-89.2 and 69.0-6.4%, resp., while in 1410 min. it gave 188.8-204.4 and 80.1-79.0%, resp.; IX in 5 min. gave 109.7-5.7 and 82.5-72.3%, resp.; IV in 5 min. gave 30.6-8.3 and 40.2-2.0%, resp. The bromination expts. gave the following nos. of substitution and addn., resp.: I 17-32 and 194-6; II 21-7 and 185-90; III 49-7 and 157-61; IV 17-14 and 162-55; XI — and 161; XII 6-9 and 135-6; VI 11-10 and 249-6; X 33-19 and 114-16; VII 179-87 and 119-14; VIII 312-36 and 19-18; IX 223-13 and 50-64. The yields of Grignard reactions with the above ketones, cited from previous papers (references given above), are also affected by the extent of conjugation, but the results must be interpreted with care since 1,4-addn. is also possible, although the generally higher yields are assumed, with systems having a

low degree of conjugation, with allowances being made for the ease of enolization. V. V. I. Ruzov and V. V. Sergovskaya. *Ibid.* 1730 3. — lithyldesacetone (I) has the highest degree of conjugation of unsats. in the RCH:CHAc series, i.e., increase of R increases the independence of the ethylene bond. I does not react in aq. alc. soln. with 0.1 N iodine, while with 0.2 N soln. the iodine no. is but 1.57% of theory, and even a 0.5 N soln. gives but 2.39% iodine no. (of theory); the butylidene analog gives 1.53, 4.39, and 17.75% of the theoretical iodine no. under the same conditions. Best prepn. of I: 550 ml. Me₂CO and 700 ml. 1.4% aq. NaOH treated at 0° with 120 g. AcH in 280 ml. H₂O, precooled to -2° over 6 hrs., let stand overnight at 0°, neutralized with AcOH, and acid. with NaCl, gave on extra. with H₂O and distn. in the presence of a crystal of iodine, 13.6% of the product, bp 119-21°, d₄²⁰ 0.8514, n_D²⁰ 1.4339. Bromination of I - at 0° in CCl₄ gives 98% of the theoretical uptake, even with twice the theoretical amt. of Br; no HBr is evolved. (G. M. Kosolopoff)

TA 2/50130

YERAP'Y, V. I.

USSR/Chemistry - Alcohols
Synthesis

JUL 49

"The Synthesis of Tertiary Alcohols Using Camphor
as a Base, I." V. I. YERAP'Y, M. I. MOVLKOV,
Lab of Org Chem, Ural State U, Sverdlovsk, 62 pp

"Zhur Obshch Khim" Vol XIX, No 7

Yield of tertiary methylborneol, synthesized by
interaction of camphor and CH_3MgI , is decreased
at high temperatures chiefly because of enoliza-
tion of camphor. At 160°C , enolization of camphor
is accompanied by formation of secondary compounds
containing camphor, latter reaction being chief
cause of a reduced yield of tertiary alcohol

2/50130

USSR/Chemistry - Alcohols
Synthesis (Contd)

JUL 49

When synthesis is carried out at -150°C . At high
temperatures chief secondary reaction is reduc-
tion of camphor into borneol. Submitted
29 Mar 48.

2/50130

CA

Characterization of α,β -unsaturated ketones. VI.
V. I. Esafey (Ural State Univ., Sverdlovsk). *Zhur.*
Obshchaya Khim. (J. Gen. Chem.) 20, 1050-7 (1950); cf.
C.I. 44, 6810d. α,β -Unsatd. ketones with normal
structure are more readily hydrolyzed with 0.01 N than
with 0.1 N NaOH; ketones similar to mesityl oxide be-
have in the reverse manner. Such gradation of reactivity
may be used in the characterization of ketones. Ketones
analogous to trialkylvinyl ketone, i.e. $R_3C:CCOR$, are
even more resistant to hydrolysis. In the 1st series
0.1-0.2-g. samples were heated in ampuls with H_2O , 0.01
or 0.1 N NaOH (10 ml.) 2 hrs. at 100° , and the CHI_2
reaction on 10-ml. aliquots used as the relative measure of
iodine consumption, calcd. on the starting material. In a
2nd series the alk. iodine soln. was allowed to react with
the 10-ml. aliquot for 15 sec., 1, 5, or 15 min. so as to give
a measure of the abs. amts. of the reacted ketones. The
results follow. $PhCH:CHCOMe$: in H_2O 61.33% re-
acted, in 0.01 N NaOH 80.91%, in 0.1 N 73.5%. C_6H_5
 $CH:CHCOMe$, 60.3, 81.0, 71.71%, resp. Mesityl oxide,
44.68, 75.43, 80.78%. $EtCMe:CHCOEt$, 43.21, 60.16,
92.16%. $EtCMe:CMcCOMe$, 37.08, 37.40, 46.69%.
The ketones with normal structure have a higher degree of
conjugation than occurs in branched-chain analogs and
this is held to be the reason for their more easy cleavage.

CA

use of the iodoform reaction in the analysis of some organic compounds. V. I. Kozlov and N. M. Staleeva (A. M. Gor'kiy Ural State Univ., Sverdlovsk, U.S.S.R.). *Zhur. Anal. Khim.* 6, 193-200 (1961). -The iodoform reaction, i.e., the reaction of compounds containing the group CH_3CO - or $\text{CH}_3\text{CH}(\text{OH})$ - with I_2 followed by hydrolysis and formation of CHI_3 , was pos. also with α,β -unsatd. ketones and their corresponding ketols not having these groups but capable of yielding acetone or satd. methyl ketones upon hydrolysis, e.g., 3-methyl-3-heptene-5-one and 4-octene-3-one. Therefore, the detn. based on CHI_3 reaction of α,β -unsatd. methylketones and of their β -ketols which upon hydrolysis yield 2 mols. of satd. methyl ketone or aldehyde and methyl ketone is inaccurate. M. Horch

ISSR/Chemistry - Diene Hydrocarbons Apr 52

"Bromination and Iodoxidation of 1,1-Dialkylbutadienes-2,3. Part VIII," V. I. Yezhov, Lab of Org Chem, Ural State U

"Zhurn Obshch Khim" Vol XXII, No 4, pp 604-611

Dehydration of dialkyl allyl carbinols in presence of iodine proceeds in 2 directions with the formation of diene hydrocarbons both with isolated bonds and with a conjugated system of double bonds. The yield of the latter is the lower, the lower the degree of hydrogenation of the alkyl group attached to the carbinol C-atoms. In 1,1-dimethylbutadiene-1,3, in

224736

contrast to all-isopropenyl and hexadiene-2,4 the double bonds are less strongly conjugated, and the bromination or iodoxidation proceeds more fully. Tertiary allyl radical and the C-atom of the carbonyl radical.

224736

YEZHOV, V.I.

YESAFOV, V.I.

**Role of Russian chemists in developing methods of synthesis by
means of zinc organic compounds. Trudy Inst. int. est. i tekhn. vol. 6:
318-346 '55. (MLWA 9:5)**

(Zinc organic compounds)

"APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962920002-0

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962920002-0"

"APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962920002-0

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962920002-0"

"APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962920002-0

APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962920002-0"

"APPROVED FOR RELEASE: 03/15/2001

CIA-RDP86-00513R001962920002-0

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
2227
2228
2229
2230
2

YESAFOV, V.I.

Chemical structure of $C_{12}H_{14}$ and synthesis of new phenylated
diene (1,3) hydrocarbons. Part 7. Zhur.ob.khim. 27 no.10:2667-2675
O '57. (MIRA 11:4)

1.Ural'skiy gosudarstvennyy universitet.
(Chemical structure) (Hydrocarbons)

AUTHOR: Yesafov, V. I.

79-20-4-17/69

TITLE: Investigation of the Thermal Decomposition of Magnesium
Iodide Ethers (Issledeniye termicheskogo razlozheniya
effiratsv vodorodnogo magniya)
I. On the Chemistry of Oxonium Compounds
(I. K khimii oksoniyevykh soedineniy)

PERIODICAL: Zhurnal Obshchey Khimii, 1958, Vol. 28, Nr 5,
pp. 1212-1218 (USSR)

ABSTRACT: The present investigation aimed at determining the
degree of the binding stability of the ether molecules of
the mentioned compounds; for this also the method of
thermal decomposition was used which has to be preferred
in the given case to other purely chemical methods. It was
found that the ethers form diethers with magnesium iodide
and this with a weak binding of the magnesium with the
oxygen of the ether molecules whereby they decompose into
thermal decomposition to magnesium iodide and ether. It
showed that the ethers with regard to their capability to
form various kinds of magnesium iodide-diethers, differ from
each other. Some ethers form one and the same magnesium iodide

Card 1/5

Investigation of the Thermal Decomposition of Magnesium Iodide Ethers. 10-28-5-17/69

I. On the Chemistry of Oxonium Compounds

diethyl, even though it was formed from magnesium iodide. The first group yields magnesium iodide ethers of two kinds. The first is obtained of magnesium iodide as such, the other from a mixture of iodine and magnesium or by displacement from the ether from $MgI_2 \cdot 2(C_2H_5)_2O$. Ethers with light primary radicals are displaced from $MgI_2 \cdot 2(C_2H_5)_2O$ by ethers with heavy primary radicals. The thermal decomposition of the magnesium iodide diethers of the second kind with aliphatic ethers leads to a formation of olefine and ether iodide with one ether molecule and iodine atom taking part in the chemical reactions. The second iodine atom remains strongly bound to the magnesium. It was further shown that in the thermal reaction of the magnesium-iodide-ethers of the second kind with ~~at~~ aromatic ethers, ketones, alkyls, phenols and resinous masses form. One molecule of the ~~fat~~ aromatic ether separates in free state. The presence of a solid chemical bond of magnesium to the oxygen

Card 2/3

Investigation of the Thermal Decomposition of Magnesium Iodide Ethers. 79-0000-11/67

I. On the Chemistry of Oxonium Compounds

of the ~~an~~ ether molecule in the magnesium iodide diether of the second kind proves its oxonium nature. Based on a scheme for the thermal decomposition of the magnesium iodide diether of the second kind was proposed. There are 6 references and 1 of which are cited.

ASSOCIATION: Ural'skiy gosudarstvennyy universitet
(Ural State University)

SUBMITTED: May 4, 1967

Card 4/4

AUTHOR: Yesafov, V. I.

79-26-5-18/69

TITLE: Investigation of the Thermal Decomposition of the Reaction Products of Dioxane With Magnesium Iodide-Diether, Magnesium-Iodide and the Mixture of Magnesium and Iodine (Izucheniye termicheskogo razlozheniya produktov vzaimodeystviya dioksana s diefiratom yodistogo magniya, yodistym magniyem i smes'yu magniya s yodom) II. On the Chemistry of Oxonium Compounds (II. K khimii oksoniyevykh soedineniy)

PERIODICAL: Zhurnal Obshchey Khimii, 1958, Vol. 28, Nr 5, pp. 1218 - 1221 (USSR)

ABSTRACT: After it was found that ethers with magnesium iodide yield compounds of the formula $MgJ_2 \cdot 2ROR$ which on heating decompose again to their initial products, it was of interest to investigate analogous compounds containing dioxane. Above all it was found that in the action of dioxane on $MgJ_2 \cdot 2(C_2H_5)_2O$ and $RMgHal \cdot 2(C_2H_5)_2O$ (Reference 2) a complete analogy exists. The dioxane displaces equimolecularly the ethyl ether from these compounds. Different from the ethers, the compound

Card 1/3

79-28-5-1E/69

Investigation of the Thermal Decomposition of the Reaction Products of Dioxane With Magnesium Iodide-Diether, Magnesium Iodide and the Mixture of Magnesium and Iodine . II. On the Chemistry of Oxonium Compounds


MgJ_2 with dioxane, $MgJ_2 \cdot 2C_4H_8O_2$, on heating separates only half of the dioxane contained in it, the other half suffering decomposition. Thus the oxygen atoms of dioxane have a greater capability to form oxonium salts than the ethers, and they yield more stable oxonium compounds. The most typical oxonium compound of dioxane is obtained in its reaction with a mixture of magnesium and iodine. This product must be regarded as oxonium salt of symmetrical (formula I) or asymmetrical structure (II). The question if these salts form at the same time, or if first salt (I) is formed which then converts to salt (II) remains unsettled. It is only clear that in salt (I) if it remains unchanged on heating the β, β' -diiodine-diether would have to be found among the decomposition products, which was not observed. The experimental results speak fully in favor of structure (II). Among the decomposition products of this salt there were found: dioxane, ethylene, ethylene iodide.

Card 2/3

79-28-5-18/69

Investigation of the Thermal Decomposition of the Reaction Products of Dioxane With Magnesium Iodide-Diether, Magnesium Iodide and the Mixture of Magnesium and Iodine. II. On the Chemistry of Oxonium Compounds

ethyl iodide, acetic anhydride, iodine and magnesium. After all the results of the reaction of dioxane with the mixture of magnesium and iodine prove that the latter as well as the ethers in this case do not react according to the scheme



as a high-melting dioxonium salt is formed here instead of the low-melting ethylene iodide. There are 2 Soviet references.

ASSOCIATION: Ural'skiy gosudarstvenny universitet (Ural' State University)

SUBMITTED: May 3, 1957

Card 3/3

5 (3)

AUTHORS:

Yesafov, V. I., Stashkov, L. I., Sirotkin, L. B.,
Suvorov, A. L., Novikov, Ye. G.

SOV/79-29-3-20/61

TITLE:


On the Characteristics of the α, β -Unsaturated Ketones. VII
(K kharakteristike α, β -nepredel'nykh ketonov. VII)

PERIODICAL:

Zhurnal obshchey khimii, 1959, Vol 29, Nr 3, pp 845-849 (USSR)

ABSTRACT:

The present paper is issued as first publication of experimental data on the hydrolytic cleavage of the aliphatic aromatic α, β -unsaturated ketones containing an aryl radical which is directly combined with the carbonyl group. Ketones of this type were obtained by dehydration of the β -ketols which had been synthesized according to the method of Grignard, V. and Colonge, I. (Ref 2).



Experiments with respect to the hydrolytic cleavage of the β -ketols were carried out as well. The data of table 2 show that the β -ketols are far more unstable than the corresponding

Card 1/2

On the Characteristics of the α,β -Unsaturated Ketones. VII

SOV/79-29-3-20/61

α,β -unsaturated ketones and prove to be more sensitive to very weak hydrolysis reagents. Besides, the behavior of the β -ketols in the hydrolysis differs from that of the α,β -unsaturated ketones by the fact that a change of the NaOH-concentration exerts a slight influence upon the cleavage intensity of the β -ketols whereas the hydrolytic cleavage of the α,β -unsaturated ketones is considerably influenced. The rate of hydrolysis of the aliphatic aromatic ketones investigated increases significantly when the NaOH concentration is increased from 0.01 to 0.1 n. 8 β -ketols hitherto unknown were synthesized and described. It was determined how far the hydrolytic cleavage of the β -ketols and at the same time that of the α,β -unsaturated ketones develops and it was proved that the latter separate but little HBr on bromination. There are 2 tables and 4 references, 2 of which are Soviet.

ASSOCIATION: Ural'skiy gosudarstvennyy universitet (Ural State University)

SUBMITTED: February 18, 1958

Card 2/2

5(3)

AUTHOR:

Yesafov, V. I.

SOV/79-29-7-5/85

TITLE:

Synthesis of the Phenylated α,β -Unsaturated Alcohols and the (1,3)-Diene Hydrocarbons.VIII (Sintez fenilirovannykh α,β -nepredel'nykh spirtov i diyenovykh (1,3) uglevodorodov.VIII)

PERIODICAL:

Zhurnal obshchey khimii, 1959, Vol 29, Nr 7, pp 2129-2132 (USSR)

ABSTRACT:

In order to solve the problems concerning the separation of HOMgX from the forming magnesium halide carbinolates (Refs 1-3), the reaction of 2,2,3-trimethyl-5-p-arylpenten-3-one-5 ($\text{Ar}=\text{C}_6\text{H}_5$, $n=\text{CH}_3$, $n=\text{C}_2\text{H}_5$), which may be present in cis- and trans-forms with RMgX (Ref 4), are described here. The experiments showed that among the three homologous ketones 2,2,3-trimethyl-5-p-tolylpenten-3-one-5 produces the best yields in tertiary alcohols. These alcohols can be dehydrated only with difficulties. In the Grignard reaction the other two ketones mainly yielded diene hydrocarbons (1,3). This particular behavior might be explained also by the influence of a spatial factor. The phenylated diene hydrocarbons (1,3) of the structure (III) and (IV) obtained show, compared to the similar aliphatic compound (V) (Ref 3), a stronger stability of the double bond

Card 1/2

Synthesis of the Phenylated α,β -Unsaturated
Alcohols and the (1,3) Diene Hydrocarbons. VIII

SOV/79-29-7-5/83

in position 3,4 and may be brominated therefore only with the separation of a high amount of HBr. The following α,β -unsaturated tertiary alcohols were newly synthesized and characterized: 2,2,3-trimethyl-5-p-tolyhexen-3-ol-5 and 2,2,3-trimethyl-5-p-tolyhepten-3-ol-5 as well as the phenylated diene hydrocarbons (1,3): 2-phenyl-4,5,5-trimethylhexadiene-1,3 and 2-p-ethyl-phenyl-4,5,5-trimethylhexadiene-1,3. It is assumed that the stability of the halogen magnesium carbinolates formed by Grignard reaction depends on α,β -unsaturated ketones. There are 1 table and 4 Soviet references.

ASSOCIATION: Ural'skiy gosudarstvennyy universitet (Ural State University)

SUBMITTED: March 17, 1958

Card 2/2

YESAFOV, V. I.

On the history of the origin of the stereochemical theory and the attitude of A.M. Butler and some of his contemporary West European chemists toward it. Trudy Inst.ist.est.i tekhn. 304-4-135-174 '60. (MIRA 13:8)

(Stereochemistry)

(Butlerov, Alexandr Mikhailovich, 1828-1886)

86499

53610

2209, 1373, 1153

S/079/60/030/011/004/026
B001/B066

AUTHORS:

Yesafov, V. I. and Yakunina, G. I.

TITLE:

Chemistry of Onium Compounds. III. Investigation of Thermal Decomposition of the Reaction Products of Tetrahydrofuran, α -Methyl Furan, Pyrrole, Thiophene With the Dietherate of Magnesium Iodide and With Magnesium Iodide

PERIODICAL:

Zhurnal obshchey khimii, 1960, Vol. 30, No. 11,
pp. 3572-3576

TEXT: V. I. Yesafov (Refs. 1,2) showed in his papers that the etherate $MgI_2 \cdot 2(C_2H_5)_2O$ is a very convenient agent for the relative estimation of the degree of aromaticity of the five-membered O-, N- and S-heterocyclic compounds. These heterocyclic compounds may be arranged in the following order on the basis of increasing difficulty in the release of the unshared electron pair of heteroatoms with formation of "onium compounds"; and for silvan, pyrrole, and thiophene with respect to the degree of stability increase of the electron sextet of heterocycles;

Card 1/4

86499

Chemistry of Onium Compounds. III. Investigation of Thermal Decomposition of the Reaction
 S/079/60/030/011/004/026
 B001/B066
 Products of Tetrahydrofuran, α -Methyl Furan, Pyrrole, Thiophene With the
 Dietherate of Magnesium Iodide and With Magnesium Iodide



This order is further confirmed by the experimental data of thermal decomposition of the reaction products of the mentioned heterocyclic compounds with anhydrous magnesium iodide. Tetrahydrofuran forms with the latter a compound which decomposes on heating by cleaving the heterocycle (Ref. 5). Silvan and pyrrole give, only on heating with magnesium iodide, compounds which decompose at high temperature, also under cleavage of the heterocycles. Also on prolonged heating, thiophene does not react with magnesium iodide. It follows from this that furan and pyrrole, as well as their compounds, maintain the "benzene-like" state of electrons in the heterocycles only at low temperatures, with increasing temperature, however, this state is disturbed, in which connection the unshared electron pairs of oxygen and nitrogen are set free, and stable onium compounds are formed with MgI_2 .

Card 2/4

86499

Chemistry of Onium Compounds, III. Investigation of Thermal Decomposition of the Reaction Products of Tetrahydrofuran, α -Methyl Furan, Pyrrole, Thiophene With the Dietherate of Magnesium Iodide and With Magnesium Iodide



which also explains the decomposition of O- and N-heterocycles on heating. It was thus shown that 1) tetrahydrofuran displaces the diethyl ether from $MgI_2 \cdot 2(C_2H_5)_2O$ to form $MgI_2 \cdot 2C_4H_8O$ which is decomposed under cleavage of a molecule of tetrahydrofuran, that 2) silvan and pyrrole displace one molecule of ether from $MgI_2 \cdot 2(C_2H_5)_2O$, and give compounds with MgI_2 which decompose under cleavage of the heterocycles, that 3) thiophene reacts neither with $MgI_2 \cdot 2(C_2H_5)_2O$ nor with MgI_2 . It is assumed that the participation of heteroatoms in the formation of heterocycles increases their capability of forming onium compounds. This assumption is supported by the fact that dioxane and tetrahydrofuran give with MgI_2 stabler compounds than simple aliphatic ethers. G. I. Kusnetsova, I. T. Bel'skiy, Card 3/4

861,99

Chemistry of Onium Compounds. III. Investigation of Thermal Decomposition of the Reaction Products of Tetrahydrofuran, α -Methyl Furan, Pyrrole, Thiophene With the Dietherate of Magnesium Iodide and With Magnesium Iodide

and S. Z. Tayts are thanked for making available the samples. There are 11 references: 5 Soviet, 4 US, 2 British, and 2 German.

ASSOCIATION: Ural'skiy gosudarstvennyy universitet (Ural State University)

SUBMITTED: August 18, 1959

Card 4/4

YESAFOV, V.I.

Characteristics of di-tertiary β -glycols. Part 1. Zhur.ob.khim.
30 no.10:3272-3275 0 '61. (MIRA 14:4)

1. Ural'skiy gosudarstvennyy universitet.
(Glycols)

YESAFOV, V. I.

35276

C/120/62/COA/006/002/026
B101/B110

15.8220

AUTHORS:

Tager, A. A., Suvorova, A. I., Goldyrev, L. N., Yesafov,
V. I., Berestova, V. L.

TITLE:

Effect of the chemical structure of the plasticizer on the
vitrification temperature of polymers. I. Plasticizing of
polystyrene with diphenic acid and naphthalic acid esters

PERIODICAL:

Vysokomolekulyarnyye soyedineniya, v. 4, no. 6, 1962,
803-808

TEXT: Thermomechanical curves were plotted for polystyrene (PSt)
plasticized with 25 mole% of: monomethyl-, monoethyl-, and monobutyl
diphenate; dimethyl-, diethyl-, ethyl-butyl-, dibutyl-, ethyl-octyl-, and
diheptyl diphenate; dimethyl, diethyl, and dibutyl naphthalate. The
synthesis of ethyl-butyl diphenate (b.p. 167-168°C/15 mm Hg, MR 91.89)
and of ethyl-octyl diphenate (MR 110.57), now produced for the first time,
will be published. The compatibility of the plasticizer with PSt was
studied on the basis of the critical mixing temperature, which lay at
100-130°C with diphenic acid monoester, below room temperature (sometimes

Card 1/2

Effect of the chemical structure ...

S/ 90/62/CO4/CO6/CO2/CO6
B101/B110

at $\sim -50^{\circ}\text{C}$) with esters of this acid, and at room temperature with naphthalates. Results: (1) The vitrification temperature, T_v , of plasticized PST drops with increasing compatibility. Pure PST had $T_v = 105^{\circ}\text{C}$, PST with monoesters had $T_v = 40-70^{\circ}\text{C}$, PST with diphenic acid diesters yielded the lowest T_v . T_v dropped with increasing length of the alkyl radical: ethyl-octyl diphenate yielded $T_v = -11^{\circ}\text{C}$; the naphthenates showed a low effect ($T_v = 9-48^{\circ}\text{C}$). (2) With increasing content of CH_2 links in the alkyl radical, T_v of diphenic acid diesters approaches a minimum at $n_{\text{CH}_2} = 10-12$, and then rises again. (3) The structure of the aromatic radical of the plasticizer affects T_v : diphenates (and phthalates) plasticize more intensively than naphthalates. There are 3 figures and 2 tables.

ASSOCIATION: Ural'skiy gosudarstvennyy universitet im. A. M. Gor'kogo
(Ural State University imeni A. M. Gor'kiy)

SUBMITTED: March 21, 1961

Card 2/2

YESAFOV, V.I.

38277

S/190/62/004/006/003/026
B101/B110

15.02.20
AUTHORS:

Tager, A. A., Suvorova, A. I., Goldyrev, L. N., Yesafov, V. I.,
Topina, L. P.

TITLE:

Effect of the chemical structure and the size of the plasticizer molecule on the vitrification temperature of polymers. II. Plasticizing of polymethyl methacrylate with esters of diphenic and naphthalic acids

PERIODICAL: Vysokomolekulyarnyye soyedineniya, v. 4, no. 6, 1962, 809-814

TEXT: Thermomechanical curves were plotted for polymethyl methacrylate (PMMA) plasticized with 25 mole% of: monomethyl, monobethyl, and monobutyl diphenate; dimethyl, diethyl, ethyl-butyl, dibutyl, and diheptyl diphenate; dimethyl, diethyl, and dibutyl naphthalate. Results: (1) The better the compatibility between polymer and plasticizer, the greater the drop in the vitrification temperature, T_v , of pure PMMA ($T_v = 100^\circ\text{C}$).

(2) T_v dropped with increasing length of the alkyl radicals of the diphenate down to a minimum (-9°C). (3) Monoesters of diphenic acid and naphthalates showed a lower plasticizing effect ($T_v \sim 50^\circ\text{C}$). (4) The
Card 1/2

Effect of the chemical structure ...

S/19C/62/004/006/003/026
B101/B110

structure of the aromatic radical affects the plasticizing effect. The better plasticizing of diphenates is explained by the ability of the compound to be turned round the C-C bond between the two benzene rings. In the case of monoesters, the free COOH reduces the compatibility. (5) The molar concentration rule does not apply to the polymer plasticizer systems investigated. There are 5 figures and 1 table.

ASSOCIATION: Ural'skiy gosudarstvennyy universitet im. A. M. Gor'kogo
(Ural State University imeni A. M. Gor'kiy)

SUBMITTED: March 21, 1961

Card 2/2

YESAFOV, V.I.

History of the discovery of organic reactions in the presence
of anhydrous aluminum halides. Trudy Inst.ist.est.i tekhn.
39:104-140 '62. (MIRA 16:2)
(Friedel-Crafts reaction)

YESAFOV, V.I.; SHITOV, G.P.

Characteristics of primary-tertiary β -glycols. Part 3. Zhur.ob.-
khim. 32 no.9:2819-2822 S '62. (MIRA 15:9)

1. Ural'skiy gosudarstvennyy universitet.
(Glycols)

YESAFOV, V.I.; ZHUKOVA, L.P.

Characteristics of secondary-tertiary β -glycols. Part 2.
Zhur.ob.khim. 32 no.9:2816-2819 S '62. (MIRA 15:9)

1. Ural'skiy gosudarstvennyy universitet.
(Glycols)

YESAFOV, V.I.; DASHKO, V.K.; MAREK, E.M.

Characteristics of secondary-tertiary β -glycols. Part 5. Cont.
b. khim. 34 no.12:4094-4096 D '64 (MIRA 18:1)

1. Ural'skiy gosudarstvennyy universitet.

USSR / Human and Animal Physiology (Normal and Pathological).
Lymph Circulation.

T

Abs Jour : Ref Zhur - Biologiya, No 13, 1958, No. 60370

Author : Yesakov, A. I.

Inst : ~~Not given~~

Title : Investigation of the "Automatic" Action of the Center
of Lymphatic Hearts

Orig Pub : Zh. obshch. biologii, 1957, 18, No 3, 185-193

Abstract : The contraction of the posterior lymph hearts (LH) in a
frog was recorded on a kymograph. The warming of the
spinal cord with a thermode on the level of the III
vertebrum produced an increase in the contraction rate;
the destruction of the cord on that level did not
interrupt the rhythmical action of LH. It was stopped
only when the spinal cord was destroyed at the VII
vertebrum. The rhythmic activity of LH originates in the

Card 1/2

YASAKOV, A.I., Cand Bio Sci—(disc) "Study of the processes ~~in the~~ ^{in the}
~~the~~ the so-called automatic activity of lymphatic hearts." Mos, 1958.
14 pp (Mos Order of Lenin and Order of Labor Red Banner State Univ
L.V.Lomonosov), 150 copies

-23-

YESAKOV, A.I.

Metabolic factors of automatism (exemplified by the cerebrospinal center of lymph hearts). Zmr.ob.biol. 20 no.1:28-34 Ja-F '59.
(MIRA 12:2)

1. Kafedra fiziologii zhivotnykh Moskovskogo gosudarstvennogo universiteta im. M.V.Lomonosova.
(LYMPHATICS) (SPINAL CORD) (PHYSIOLOGICAL CHEMISTRY)

SNYAKIN, P.G., prof.; YESAKOV, A.I., kand.biolog.nauk

Muscular sense. Zdorov'e 7 no.9:9-10 S '61.
(MUSCULAR SENSE)

(MIRA 14:9)

YESAKOV, A.I.

Metabolic nature of the automatism of nerve cells as exemplified
in the spinal center of the lymph heart. Zhur. ob. biol. 22 no.2:
136-143 Mr-Apr '61. (MIRA 14:5)

1. Department of Animal Physiology, State University of Moscow.
(NERVES)

YESAKOV, A.I.

Problems of the efferent regulation of receptors; based on an example of lingual chemoreceptors. Biul. eksp. biol. i med. 51 no.3: 3-8 Mr '61.
(MIRA 14:5)

1. Iz laboratorii fiziologii i patologii organov chuvstv (zav. - prof. P.G.Snyakin) Instituta normal'noy i patologicheskoy fiziologii (dir. - deystvitel'nyy chlen AMN SSSR V.V.Parin) AMN SSSR, Moskva. Predstavlena deystvitel'stva chlenom AMN SSSR P.K.Anokhinym.
(RECEPTORS (NEUROLOGY)) (STOMACH)
(TONGUE)

YESAKOV, A. I.

"Methodology of neurodynamic investigations and practical work
in the physiology of human analysts" by P. O. Makarov. Reviewed
by A. I. Esakov. Nauch. dokl. vys. shkoly; biol. nauki no.3:
208-209 '62. (MIRA 15:7)

(SENSES AND SENSATION) (PHYSIOLOGY--LABORATORY MANUALS)
(MAKAROV, P. O.)

SNYAKIN, P. G., prof.; YESAKOV, A. I., kand. biologicheskikh nauk

Our habits are stereotypes. Zdorov'e 8 no.11:4-5 II '62.
(MIRA 15:10)

(HABIT)

YESAKOV, A.I.

Effect of the sympathetic nervous system on the electric activity of tongue receptors. Trudy Inst. norm. i pat. fiziologii. AMN SSSR 6:63-65 '62 (MIRA 17:1)

1. Laboratoriya fiziologii i patologii organov chuvstva (zav.-prof. P.G. Snyakin) Instituta normal'noy i patologicheskoy fiziologii AMN SSSR.

YESAKOV, A.I.; ZAYKO, N.S.

Effect of guanidine on the functional activity of taste
receptors. Fiziol. zhur. 49 no.8:984-989 Ag '63.

(MIRA 17:2)

1. From the Laboratory for Physiology and Pathology of Sense
Organs, Institute of Normal and Pathologic Physiology,
U.S.S.R. Academy of Medical Sciences, Moscow.

YESAKOV, A.I.

Reflex regulation of "spontaneous" activity of tongue chemo-receptors. Biul. eksp. biol. i med. 56 no.8:7-11 Ag '63.

(MIRA 17:7)

1. Iz laboratorii fiziologii i patologii organov chuvstv (zav. - prof. P.G. Snyakin) Instituta normal'noy i patologicheskoy fiziologii (dir. - deystvitel'nyy chlen AMN SSSR prof. V.V. Parin) AMN SSSR, Moskva. Predstavleno deystvitel'nyy chlenom AMN SSSR P.K. Anokhinym.

YESAKOV, A.I.

Efferent reactions in the hypoglossal nerve. Biul. eksp. biol. i med.
56 no.11:15-18 0 [i.e. N] '63. (MIRA 17:11)

1. Iz laboratorii fiziologii i patologii organov chuvstv (zav. - prof.
P.G. Snyakin) Instituta normal'noy i patologicheskoy fiziologii (dir.-
deystvitel'nyy chlen AMN SSSR prof. V.V. Parin) AMN SSSR, Moskva.

YESAKOV, A.I.

Electrophysiological analysis of the functional mobility and of processes regulating the taste receptor apparatus. Trudy Inst. norm. i pat. fiziol. AMN SSSR 7:43-44 '64. (MIRA 18:6)

1. Laboratoriya fiziologii i patologii organov chuvstv (zav. - prof. P.G. Snyakin) Instituta normal'noy i patologicheskoy fiziologii AMN SSSR.

YESAKOV, A.I. (Moskva)

Role of different regulation levels of receptors in the
perception process. Vest. AMN SSSR 21 no.1:62-68 '66.

(MIRA 19:1)

YESAKOV, A.I.; FILIN, V.A.

Physiological characteristics of the functioning of the taste
receptor apparatus. Fiziol. zhur. 50 no.2:169-176 F '64.

(MIRA 18:2)

1. Laboratoriya fiziologii i patologii organov chuvstv Instituta
normal'noy i patologicheskoy fiziologii AMN SSSR, Moskva.

YESAKOV, A.I.

Possible specific stimulus of the "automatic" center of the lymph heart. *Biul. eksp. biol. i med.* 57 no.6:19-22 Je '64. (MIRA 18:4)

1. Kafedra fiziologii zhivotnykh Moskovskogo gosudarstvennogo universiteta imeni Lomonosova.

YESAKOV, I.S.; YASHCHENKO, Z.G.

Interpreting vertical electric sounding curves by the T method.
Razved. i prom. geofiz. no.30:50-54 '59. (MIRA 12:12)
(Electric prospecting)

YASHCHENKO, Z.G.; YESAKOV, I.S.

Use of electric prospecting in studying the elastic properties
of igneous rocks. Razved. i prom. geofiz. no. 33:23-29 '59.
(MIRA 13:4)

(Rocks--Electric properties) (Elasticity)

YESAKOV, V.A.

M.S. Bodnarskii; obituary. Izv. AN SSSR Ser. geog. no. 1:95-96
Ja-F '54. (MLRA 7:2)
(Bodnarskii, Mitrofan Stepanovich, 1870-1953)

USSR/Geography Education

Card : 1/1 Pub. 45 - 8/20

Authors : Esakov, V. A.

Title : History of the origination of geography faculties at Russian universities

Periodical : Izv. AN SSSR. Ser. Geog. 4, 51 - 60, July - August 1954

Abstract : History of the establishment of geography faculties at Russian universities since the second half of the 19th century.

Institution : Acad. of Sc. USSR, Institute of Natural History and Technique

Submitted :

YESAKOV, V. A.
USSR/Geophysics - Moscow University

FD-1152

Card 1/1 Pub. 129-16/23

Author : Yesakov, V. A.

Title : From the history of geography in Moscow University (up to the establishment of the chair of geography in 1884)

Periodical : Vest. Mosk. un., Ser. fizikomat. i yest. nauk, 9, No 7, 131-136, Oct 1954

Abstract : The Petersburg Academy of Sciences, founded in 1725, established in 1739 its Geographical Department. The author lists the numerous studies and expeditions undertaken after this date.

Institution :

Submitted : May 22, 1954

YESAKOV, V.A.; SOLOV'YEV, A.I., redaktor; VOLODINA, N.I., redaktor;
~~LEVINAKOVA~~, T.A., tekhnicheskii redaktor

[D.N. Anuchin and foundation of the Russian academic geographical school.] D.N. Anuchin i sozдание russkoi universitetskoi geografi-
cheskoi shkoly. Moskva, Izd-vo Akademii nauk SSSR, 1955. 180 p.
(Anuchin, Dmitrii Nikolaevich, 1843-1923) (MIRA 8:10)

YESAKOV, V.A.

M.I. Veniukov's correspondence with N.M. Przheval'skii.

Vop. ist.est. i tekhn. no.1:207-212 '56.

(MLRA 9:10)

(Veniukov, Mikhail Ivanovich, 1832-1901)

(Przheval'skii, N.M.)

3(0)

SOV/10-59-3-21/32

AUTHOR: Yesakov, V.A.

TITLE: Scientific Contacts of A. Humboldt with Russian Scientists

PERIODICAL: Izvestiya Akademii nauk SSSR, Seriya geograficheskaya, 1959,
Nr 3, pp 124-130 (USSR)

ABSTRACT: This is a review of scientific contacts of A. Humboldt with
Russian scientists. There is 1 photograph, and 18 references,
of which 15 are Russian and 3 German.

ASSOCIATION: Institut istorii yestestvoznaniya i tekhniki AN SSSR (the
Historical Institute of the Science of Natural History and Tech-
nology, AS USSR).

Card 1/1

3(

SOV/10-59-4-28/29

AUTHOR: Yesakov, V.A.

TITLE: A. Humboldt's Anniversary in the USSR

PERIODICAL: Izvestiya Akademii nauk SSSR, Seriya geograficheskaya
1959, Nr 4, p 158 (USSR)

ABSTRACT: The article is concerned with the 100th anniversary of the death of A.v. Humboldt in the USSR. The following personalities and organizations participated in the commemoration of the great German scientist: Academician D.I. Shcherbakov, Corresponding Member AN SSSR (AS USSR), S.V. Kalesnik, Academician A.A. Grigor'yev, Corresponding Member AS USSR S.V. Obruchev, the Akademiya nauk SSSR (Academy of Sciences USSR), the Geograficheskoye obshchestvo SSSR (Geographical Society USSR), the Glavnaya geofizicheskaya observatoriya im. A.I. Voyeykova (Central Geophysical Observatory imeni A.I. Voyeykov), the Astronomicheskii institut imeni P.K. Shternberga (Institute of

Card 1/3

SOV/10-59-4-28/29

A. Humboldt's Anniversary in the USSR

Astronomy imeni P.K. Shternberg), the Vsesoyuznoye botanicheskoye obshchestvo (All-Union Botanical Society), the Otdeleniye geologo-geograficheskikh nauk (Section of Geological and Geographical Sciences) AS USSR, the Institut istorii yestestvoznaniya i tekhniki (Institute of History of the Natural Sciences and Technology), the Sovetskoye natsional'noye ob'yedineniye istorikov yestestvoznaniya i tekhniki (Soviet National Association of Historians of Natural Sciences and Technology), the Moskovskiy gosudarstvennyy universitet im. M.V. Lomonosova (Moscow State University imeni M.V. Lomonosov), the Moskovskoye obshchestvo ispytateley prirody (Moscow Society of Naturalists), Obshchestvo sovetsko-germanskoy družby i nauchnykh svyazey (Soviet-German Society for the Promotion of Friendship and Scientific Connections), the Gosudarstvennyy istoricheskiy muzey (State Historical Museum), the Gosudarstvennyy muzey

Card 2/3

SOT/10-59-4-28/29

A. Humboldt's Anniversary in the USSR

izobrazitel'nykh iskusstv (State Museum of Graphic Arts), the Muzey zemlevedeniya MGU (Geographical Museum MGU), the Gosudarstvennoye izdatel'stvo geograficheskoy literatury (State Publishing House of Geographical Publications) and the Ministerstvo svyazi SSSR (Ministry of Communications USSR). There is 1 Soviet reference.

Card 3/3

YESAKOV, V.A.; SOLOV'YEV, A.I.; FEDOSEYEV, I.A., otv. red.;

[Russian geographical explorations of European Russia and
the Urals in the 19th and the beginning of the 20th century]
Russkie geograficheskie issledovaniia Evropeiskoi Rossii i
Urala v XIX - nachale XX v. Moskva, Nauka, 1964. 177 p.
(MIRA 17:11)

NAUMOV, Guriy Vasil'yevich; FEDOSEYEV, I.A., otv. red.; YESAKOV,
V.A., red.; SOLOV'YEV, A.I., red.

[Russian geographical explorations in Siberia in the 19th
century] Russkie geograficheskie issledovaniia Sibiri v
XIX - nachale XX v. Moskva, Nauka, 1965. 146 p.
(MIRA 19:1)

YESAKOV, V.A.; PLAKHOTNIK, A.F.; ALEKSEYEV, A.I.; FEDOSEYEV, I.A.,
otv. red.; SOLOV'YEV, A.I., red.

[Russian ocean and sea studies in the 19th to the beginning of the 20th century] Russkie okeanicheskie i morskoe issledovaniia v XIX-nachale XX v. Moskva, Nauka, 1964.
158 p. (MIRA 18:1)

YESAKOV, Vasilii Petrovich; PARFENOV, Eduard Yevgen'yevich;
PROZOROV, Valentin Alekseyevich; LERNER, D.M., red.

[Automated electric drive systems with regulated semi-
conductor rectifiers] Sistemy avtomatizirovannogo elektro-
privoda s upravliaemymi poluprovodnikovymi vypriamiteliami.
Leningrad, 1964. 35 p. (MIRA 17:11)

YESAKOV, V.P., inzhener.

Operating conditions of electric winches on ships. Sudostroenie
23 no.3:34-35 Mr '57. (MLRA 10:5)
(Winches) (Electricity on ships)

YESAKOV, V.P., kand.tekhn.nauk

Systems for electric propelling units using steady current. Sudostroenie
25 no.2:31-34 F '59. (MIRA 12:4)
(Electricity on ships)

L 33115-66

ACC NR: AP6024083

SOURCE CODE: UR/0144/66/000/002/0235/0236

AUTHOR: Zav'yalov, A. S.; Got'man, A. A.; Molchanov, V. D.; Krasnyuk, N. P.;
Agranovskiy, K. Yu.; Berger, A. Ya.; Greyer, L. K.; Yezakov, V. P.; Miller, Ye. V.;
Pyatman, K. I.; Abryutin, V. N.; Gubanov, V. V.; Oranskij, M. I.; Yevseyev, M. Ye.;
Morkin, G. B.; Sinol'nikov, Ye. M.; Avilov-Karnaukhov, B. N.; Bogush, A. G.;
Bolyayov, I. P.; Pekker, I. I.; Chernyavskiy, F. I.

ORG: none

TITLE: O. B. Bron (on his 70th birthday)

SOURCE: IVUZ. Elektromekhanika, no. 2, 1966, 235-236

TOPIC TAGS: electric engineering personnel, circuit breaker

ABSTRACT: Osip Borisovich Bron was born in 1896 in Klintsi. In 1920, he graduated from the physics-math faculty of Khar'kov Technological Institute. He became a professor in 1930. He defended his doctor's thesis in 1940. During the second world war, he was in the navy. After demobilization in 1950, Engineer Colonel Bron went to work teaching at the Leningrad Industrial Correspondence School. He became the head of the Chair of Theoretical Bases of Electrical Technology in 1958. He is closely associated with scientific and development work, and has cooperated closely in this area with the Leningrad "Elektrosila" plant since 1946. His work has been in the areas of spark-damping and high-power circuit breakers. He has published over 140 scientific works and 19 inventions. [JPRS]

SUB CODE: 05, 09 / SUBM DATE: none

Card 1/1

5.3300

77094

307/62-57-12-38/43

AUTHORS: Zhuze, T. P., Zhurba, A. S., Yesakov, Ye. A.

TITLE: Brief Communications. Investigation of P-V-t-N Relation and Phase Equilibrium in Ethylene-Cyclohexane System

PERIODICAL: Izvestiya Akademii nauk. Otdeleniye khimicheskikh nauk, 1959, Nr 12, pp 2251-2253 (USSR)

ABSTRACT: Investigation of binary systems of unsaturated gases and paraffins, naphthenes, and aromatic hydrocarbons, presents great practical interest in view of the high solubility of the latter in compressed unsaturated gases. P-V-t-N relationship of the ethylene-cyclohexane system was studied by the authors in a modified apparatus described by Sage and Lacey (Trans. Amer. Inst. Mining Met. Engrs., 1940, Nr 136, p 138). Isotherms $V = f(p)t$ were traced in the range from 30 to 150° for ethylene-cyclohexane mixtures with 20 to 85 molar % ethylene, at pressures ranging from 10 atm to pressures somewhat above the saturation point of each mixture. Saturation pressures and the corresponding specific volumes were determined from these isotherms, and dew point pressures were established in a series of separate experiments.

Card 1/3 2

Brief Communications. Investigation of
P-V-t-N Relation and Phase Equilibrium
in Ethylene-Cyclohexane System

77094

SOV/62-59-12-38/83

Data thus obtained, served to trace isotherms
 $p = f(N)t$ (where N is the molar share of ethylene
dissolved in cyclohexane); isotherms of the equilibrium
constant for ethylene and cyclohexane, $K = f(P)t$; and
isotherms of molar volumes V_M of the binary system
plotted against molar share N_2 of ethylene at 50 atm.

In this manner, the composition of the coexisting
phases, the equilibrium constants of ethylene and
cyclohexane at pressures up to 100 atm, and the molar
volumes of mixtures at their saturation pressures,
were determined in the temperature range from 30 to
125°. Molar volume isotherms at low temperatures
were practically linear up to $N_2 = 0.65$; at higher
temperatures, the molar volume increases sharply
starting with $N_2 = 0.50$. There are 3 figures; and
3 references, 1^{U.S.}, 2 Soviet. U.S. reference is:
B. H. Sage, W. N. Lacey, Trans. Amer. Inst. Mining.
Met Engrs., 136, 138 (1940).

Card 2/82

Inst. Geology, Processing Thermal Fuels

USSR USSR

ZHUZE, T.P.; YUSHKEVICH, G.N.; USHAKOVA, G.S.; TESAKOV, Ye.A.

Critical parameters for oil and oil-gas systems. Neft.
khoz. 41 no.6:25-31 Je '63. (MIRA 17:6)

AFANAS'YEVA, V. B.; YESAKOVA, N. P.

Relation between the snow cover and G. J. Wangenheim's types
of circulation. Trudy GGO no.151:77-80 '64. (MIRA 17:7)

ACCESSION NR: AT4046059

S/2531/64/000/166/0182/0188

AUTHOR: Yudin, M.I.(Doctor of physico-mathematical sciences); Yesakova, N. P.; Afanas'yeva, V. B.

TITLE: Preliminary evaluation of the prognostic significance of the information obtained from meteorological satellites

SOURCE: Leningrad. Glavnaya geofizicheskaya observatoriya. Trudy*, no. 166, 1964. Voprosy* interpretatsii dannykh meteorologicheskikh sputnikov (Problems in the interpretation of data of meteorological satellites), 182-188

TOPIC TAGS: meteorology, meteorological satellite, cloud, precipitation, weather forecasting, long-range weather forecasting, snow cover, radiation balance

ABSTRACT: The objective of this paper was to develop a method for the preliminary characterization of anomalies of cloud cover, the radiation balance of the underlying surface and the limits of snow and ice cover for subsequent use of such characteristics in long-range weather forecasting. The authors establish statistical relationships between such anomalies and the characteristics of future weather (temperature and precipitation); certain direct characteristics of atmospheric circulation are also analyzed in relation to future weather. Determination of the characteristics of anomalies of the cloud cover,

Card 1/3

ACCESSION NR: AT4046059

the boundaries of the snow and ice cover and the radiation balance was done using mean 10-day values for the period September-November 1948-1957. These values were mapped, after which the parameters characterizing the fields of individual elements were determined. The method used for constructing the maps and defining the characteristics of anomalies is described briefly. The state of atmospheric circulation was described using the zonal index devised by Ye. N. Blinova, the M.I. Yudin meridional index and the A. A. Rozhdestvenskiy hydrodynamic indices. These parameters were used to supplement the 10-day means of temperature and precipitation for an analysis of these values determined for a grid of points covering much of the European SSSR. Synchronous statistical relationships were established between the 10 mentioned parameters; asynchronous prognostic relationships also were determined. The ten considered parameters were correlated with temperature and precipitation for the 10 days which followed. The computations of the correlation coefficients were performed on a "Ural-1" electronic computer. Most of the results of the computations were plotted on maps, and 66 such maps were constructed. In a considerable number of cases relationships were discovered which are characterized by quite high correlation coefficients and with a stable identical

ACCESSION NR: AT4046059

sign for the entire considered area. It was found that the selected parameters generally give more information for prediction of temperature than for prediction of precipitation. However, an absence of prognostic relationships is noted on a number of maps. The method described made it possible to establish a number of parameters of the state of the atmosphere and the underlying surface which are quite closely related to the characteristic of future weather for 10 days in advance. The greater part of the parameters apply to those elements which cannot be determined globally except by use of meteorological satellites. This emphasizes the great importance of satellite observations for long-range forecasting. Orig. art. has: 2 formulas, 5 figures and 1 table.

ASSOCIATION: Glavnaya geofizicheskaya observatoriya Leningrad (Main Geophysical Observatory)

SUBMITTED: 00

ENCL: 00

SUB CODE: ES

NO REF SOV: 006

OTHER: 000

Card 3/3

AFANAS'YEVA, V.B.; YESAKOVA, N.P.

Statistical relations between the anomalies of certain
weather characteristics. Trudy GGO no.165:105-113 '64.
(MIRA 17:9)

AFANAS'YEVA, V.B.; YESAKOVA, N.P.

Relation of the shift of the boundary of snow cover to the G.I.A.
Vangengsim types of circulation in the spring period. Trudy GGO
no.168:45-48 '65.

YULIN, M.I.; YESAKOVA, D.P.; AFANAS'YEVA, V.B.

Preliminary evaluation of the prognostic significance of
information returned by meteorological satellites. Trudy
GGO no.166:182-188 '64.

(MIRA 17:11)

YESAKOVA, R.

Objective method for determination of the liming stage of gelatin stock. R. Gorodetskaya, M. Sheremet, M. Shakhnazarova, D. Virnik, V. Smirnova, and R. Yesakova. Myasnaya Ind. S. S. S.R. 25, No. 5, 52-4(1954). —The procedure for detg. the status of the liming of gelatin stock is based on extg. a sample and detg. extd. gelatin colorimetrically by means of the biuret reaction. Results are given for extractable gelatin in bone stock at 5-day intervals for 40 days of liming. Total extractable gelatin is detd. for various bones and other gelatin stock.

M. M. Fiskur

GORODETSKAYA, R.V., kandidat khimicheskikh nauk; SHAKHNazarova, M.Sh.,
mladshiy nauchnyy sotrudnik; SHEREMET, M.V.; VIRNIK, D.I.;
SMIRNOVA, V.Ye.; YESAKOVA, R.

Reducing losses in gelatin production. Trudy VNIIMP no.7:108-113
'55. (MLRA 9:8)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut myasnoy promyshlen-
nosti (for Gorodetskaya, Shakhnazarova, Sheremet); 2. Moskovskiy
zhelatinovyy zavod (for Virnik, Smirnova, Yesakova).
(Gelatin)

GORODETSKAYA, R.V., kandidat khimicheskikh nauk; SHAKHNAZAROVA, M.Sh.,
mladshiy nauchnyy sotrudnik; SHERMET, M.V.; VIRNIK, D.I.;
SMIRNOVA, V.Ye.; YESAKOVA, R.

Methods of determining the degree of liming in gelatigenous tissues.
Trudy VNIIMP no.7:114-122 '55. (MLRA 9:8)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut myasnoy promy-
shlennosti (for Gorodetskaya, Shakhnazarova, Sheremet); 2. Moskov-
skiy zhelatinovyy zavod (for Virnik, Smirnova, Yesakova).
(Gelating)